

### Familial Polycythemia Vera in Father and Daughter

*To the Editor:* Polycythemia vera (PV) is a chronic myeloproliferative disorder which usually affects the elderly. We report a familial case of PV involving a father and daughter; the latter patient was diagnosed at the relatively young age of 24 years. Human leukocyte antigen (HLA) typing revealed a possible HLA haplotype associated with familial PV.

#### PATIENT 1

A 57-year-old Japanese man was admitted to our hospital because of a headache in December, 1989. Laboratory findings on admission were as follows: red blood cells (RBC)  $7.14 \times 10^{12}/l$ , hemoglobin 16.1 g/dl, platelets,  $1,359 \times 10^9/l$ , and white blood cells (WBC)  $20 \times 10^9/l$ . Leukocyte alkaline phosphatase (LAP) score was elevated at 372 (normal range: 170–285). Bone marrow showed normocellularity with normal differentiation counts. Cytogenetic study demonstrated 46XY. RBC volume was 38.7 ml/kg (normal range: 25–35 ml/kg), and the arterial oxygen saturation ( $\text{SaO}_2$ ) did not suggest hypoxemia. Serum vitamin  $\text{B}_{12}$  and erythropoietin concentrations were 711 pg/ml (normal range: 230–800 pg/ml) and 6.8 mU/ml (normal range: 8–36 mU/ml), respectively. Mild splenomegaly was observed. From these findings, he was diagnosed with PV. He has been treated with ranimustine (50–100 mg/body weight, intravenously, every fourth month) with neither any side effects nor observable symptoms due to PV.

#### PATIENT 2


A 24-year-old Japanese woman, the daughter of patient 1, demonstrated abnormal blood counts on a routine screening examination in October, 1991. Laboratory findings at that time were as follows: RBC  $8.45 \times 10^{12}/l$ , hemoglobin 20.5 g/dl, platelets  $741 \times 10^9/l$ , and WBC  $10 \times 10^9/l$ . LAP score was elevated at 387. Bone marrow showed normocellularity with normal differentiation counts. Cytogenetic study demonstrated 46XX. RBC volume was 59.0 ml/kg, and  $\text{SaO}_2$  did not suggest hypoxemia. Serum vitamin  $\text{B}_{12}$  and erythropoietin concentrations were 759 pg/ml and 6.8 mU/ml, respectively. Mild splenomegaly was observed. From these findings, the patient was diagnosed with PV. She has been treated with phlebotomy without severe symptoms.

#### DISCUSSION

Figure 1 shows the results of HLA typings performed on this family. Among these family members, only patients 1 and 2 shared the haplotype of A2/B35/Cw3/DR8/DQ1. No other members demonstrated abnormal hematologic findings or shared the above haplotype, although the HLA type was not examined in one individual.

Several reports have suggested a genetic role in the etiology of PV that occurs in families [1,2]. PV appears more commonly in Jews as compared with Caucasians [3]. Regarding HLA antigens, HLA-B52 and HLA-DR7 have been found more frequently in PV patients as compared with normal healthy individuals, whereas HLA-DR4 has been documented less frequently [4]. Alternatively, some cases of familial PV appear to result from frequent exposure to organic solvents [5]. Therefore, genetic and/or environmental factors might be implicated in this disorder.

Neither of our patients was Jewish; nor had they been frequently exposed to organic solvents. They also did not possess either the HLA-B52 or HLA-DR7 antigen. Therefore, the shared haplotype of A2/B35/Cw3/DR8/DQ1 may have been responsible for this familial occurrence of PV. Further examination is necessary to clarify the existence of such a causal relationship.



HLA haplotype a : A2 B35 Cw3 DR8 DQ1  
 haplotype b : A2 B44 CwBL DR13 DQ1  
 haplotype c : A24 B55 Cw3 DR2 DQ1  
 haplotype d : A24 B7 Cw7 DR14 DQ1

Fig. 1. HLA types of family members.

TOHRU INABA  
 CHIHIRO SHIMAZAKI  
 HIDEYO HIRAI  
 TAKEHISA KIKUTA  
 TOSHIYA SUMIKUMA  
 YOSHIKAZU SUDO  
 NOBORU YAMAGATA  
 EISHI ASHIHARA  
 HIDEO GOTO  
 NAOHISA FUJITA  
 MASAO NAKAGAWA

Second Department of Medicine, Kyoto Prefectural University of Medicine, Kyoto, Japan

#### REFERENCES

1. Friedland ML, Wittels EG, Robinson RJ: Polycythemia vera in identical twins. *Am J Hematol* 10:101, 1981.
2. Miller RL, Purvis JD III, Weick JK: Familial polycythemia vera. *Cleve Clin J Med* 56:813, 1989.
3. Modan B, Kallner H, Zemer D, Yoran C: A note on the increased risk of polycythemia vera in Jews. *Blood* 37:172, 1971.
4. Stavropoulos C, Petropoulou F, Christophoraki M, Theodosoudi E, Kandidaki E, Papanicolaou M, Mavrikakis M, Gyftaki H: Frequency of HLA-A, B and DR antigens in patients with polycythemia vera. *Acta Haematol* 86:51, 1991.
5. Ratnoff WD, Gress RE: The familial occurrence of polycythemia vera: Report of a father and son, with consideration of the possible etiologic role of exposure to organic solvents, including tetrachloroethylene. *Blood* 56:233, 1980.

#### Intravenous Immune Globulins and Hypothermia

*To the Editor:* We would like to report on a patient who presented recurrent episodes of hypothermia following intravenous immune globulin (IVIG) infusions. This 58-year-old man had been diagnosed with stage IV B cell chronic lymphocytic leukemia (B-CLL) 15 years earlier. A prophylactic